

International Journal of Ethnobiology & Ethnomedicine

Review Article

A review on phytochemical and pharmacological potential of *Prosopis cineraria*

Shivali sachdeva^{1*}, Dr. Vichitra Kaushik², Dr. Vipin Saini²

¹Rayat and Bahra Institute of Pharmacy, Kharar,

²MM College of Pharmacy, Maharishi Markandeshwar University, Mullana, Ambala.

Correspondence should be addressed to **Shivali sachdeva**

Received 26 August 2014; Accepted 2 September 2014; Published 8 September 2014

Copyright: © 2014 **Shivali sachdeva** et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

In the present review, an attempt has been made to congregate the traditional, phytochemical and pharmacological studies done on important medicinal plant *Prosopis cineraria*, (Family Fabaceae). Fatty acids, tannins, alkaloids, flavonoids and glycosides were the major phytochemical compounds studied from this medicinal plant. The plant possesses the major pharmacological activities which includes analgesic, antyperlipidemic, antipyretic and antimicrobial activity. This review encompasses the potential application of the above plant in the pharmaceutical field due to its wide pharmacological activities.

Keywords: *Prosopis cineraria*, Phytoconstituents, Pharmacological activities.

Introduction

Plants and its products are used as medicine from the ancient time¹. Recently there has been a shift in universal trend from synthetic to herbal medicine². It is estimated by the World Health Organization that approximately 75-80% of the world's population uses plant medicines either partly or entirely as medicine. Interest in plant derived drug increases mainly due to the increasing use, and misuse, of existing synthetic drugs. This poses the need for search and development of new drugs to cure diseases¹.

The chemical substances of the medicinal plants which have the capacity of exerting a physiologic action on the human body are the primary features. The bioactive compounds of plants such as alkaloids, flavonoids, tannins and phenolic compounds are considered to be most important. The phytochemical research that has been done based on the ethno-pharmacological information forms the effective approach in the discovery of new medicinal agents from higher plants³

Prosopis cineraria (L.) (Family Fabaceae) is a promising multipurpose tree species. The tree,

locally called as Jandi or Khejri holds an important place in the rural economy in the northwest region of Indian subcontinent. The genus *Prosopis* comprises about 44 species distributed mainly in dry regions of Southwest Asia, Africa and, predominantly America from western North America to Patagonia⁴. There are various common names for the plant. In Hindi and Sanskrit it is known as Khejri and in Rajasthan known as Janti/Loong tree. In Punjab it is known by the name of Jand and in Gujarat is known as Sami, Sumri. Tamil and Jammi and Vanni in Telugu. In Sind it is known as Kandi⁵. It is also known as “wonder tree” and “king of desert” as all the parts of tree are useful. It is a small thorny, irregularly branched tree, 5-10 in high. It has thick, rough gray bark with deep fissures. Roots are very deep; the tap root of *P. cineraria* may penetrate vertically up to 20 m or more⁶. The leaves form good fodder for camels, goats and donkeys. The pods are used as a vegetable. The flowers are useful for honey production. Khejri is also used for soil improvement and sand dune stabilization. The wood is ideal for domestic heating. The bark of the tree has abortifacient and laxative properties. Khejri is reputed for the treatment of asthma and worm⁷.

Plant Description

Prosopis cineraria is a tree to 6.5 m high with cinereous cortex with intermodal prickles. It produces new flush leaves before summer. The flowers are small in size and yellow or creamy white in colour; appear from March to May after the new flush of leaves. The pods are formed soon thereafter and grow rapidly in size attaining full size in about two months time. *Prosopis cineraria* is a tree to 6.5 m high with cinereous cortex with intermodal prickles, scattered, straight and somewhat macroscopic and with conical broad bases.

Root: Root is a taproot more than 3 m long.

Leaves: Leaves are 1-3-jugate, glabrous or puberulous; petiole and rachis is 0.5-4 cm long, the

pinnae is 2-7 cm long; leaflets are 7-14-jugate, ovate, straight to subfalcate, without nerves (or 2-4-nerved at base, the midrib excentric), mucronate, 4-15 mm long x 2-4.5 mm broad, grayish when dry; stipules foliaceous, deciduous.

Flower: Flowers are yellow, glabrous; calyx truncate, 0.8-1.2 mm long; corolla 3.5 mm long, glabrous, the petals rolled back in age; anthers 0.8-1 mm long; pistil glabrous.

Fruit: Fruit is slender, elongate, 8-19 cm long (including the stipe 0.8-2 cm), subcylindrical, 4-7 mm in diameter, glabrous; pericarp is thin, brittle; endocarp segments are thin, longitudinal, little developed.

Seeds: Seeds are distant, longitudinal, ovate, 6 mm long, the tegument with open horse-shoe fissural line on faces, 10-15 in a pod, brown⁸.

Phytochemical Review

Literature survey of *Prosopis cineraria* revealed the presence of alkaloids, fatty acids, glycosides and sterols whereas glucosides are reported from flowers and flavones from seeds⁹⁻¹⁰.

Phytochemical investigations on the leaves of the plant showed the presence of hydrocarbons and phenolic acid derivatives^{11, 12}. The leaves also showed the presence of large proportion of unsaturated fatty acids, with linoleic acid and oleic acid¹³.

Phytochemical screening of the dried unripe pods of *Prosopis cineraria* showed the presence tannins, alkaloids, flavonoids and glycosides in the dried unripe pods¹⁴.

Numerous bioactive compounds such as flavonoids, alkaloids, diketones, phenolic contents, free amino acids, patulitrin, spicigerin, prosogerin A,B,C,D, lipids, b-sitosterol, sugars and vitamins have been isolated from various parts of the plant^{15,16}. The constituents and the part of the plant in which they are present have been shown in table No 1.

Table No 1

Part of the Plant	Chemical Constituent
Whole Plant	Alkaloids, fatty acids, glycosides and sterols
Leaves	Phenolic Acid derivatives, unsaturated fatty acids, linoleic acid and oleic acid.
Pods	Tannins, alkaloids, flavonoids and glycosides.
Flowers	Glucosides
Seeds	Flavones

Pharmacological Review

Ancient literature has reported the use of *P. cineraria* as folk medicine for various ailments. Its flowers mixed with sugar when administered orally prevent miscarriage¹⁷. With twig, the flowers are also known as an anti-diabetic agent. Dry pods of the plant help in preventing protein calorie malnutrition and iron calcium deficiency in blood. Smoke of leaves is used to cure eye infections¹⁸. Bark of the tree is used in the treatment of asthma, bronchitis, dysentery, leucoderma, leprosy, muscle tremors and piles^{19, 20}. Pharmacological activities like analgesic, antipyretic, antihyperglycemic, antioxidant, antihypercholesterolemic, antitumor, nootropic have been reported from different plant extracts.

Analgesic and antipyretic activity

The Petroleum ether extract of stem bark exhibited a significant antipyretic activity using Brewer's yeast induced hyperpyrexia model in experimental rats. The ethanolic extract of root was evaluated by using tail immersion and hot plate method and showed significant results. The aqueous extract of leaves was evaluated for analgesic activity by using acetic acid induced writhing test model. The Analgesic activity exhibited in Swiss Albino mice was significant as compared to control. The extract also exhibited a significant antipyretic activity at same dose using Brewer's yeast induced hyperpyrexia model²¹.

Anti hyperglycemic and antioxidant activity

The stem bark was evaluated for anti hyperglycemic activity using Alloxan induced

Hyperglycemia Model. Declined activity of antioxidant enzymes and concentration of non-enzymatic antioxidants were also normalized by drug treatment, thereby reducing the oxidative damage in the tissues of diabetic animals and hence indicating anti-diabetic and antioxidant efficacy of the extract²².

Antitumor activity

Hydro alcoholic extract of Leaves and bark were evaluated for antitumor activity against Ehrlich ascites carcinoma tumor model. Both the extracts showed significant antitumor activity²³. Methanolic extract of leaves was evaluated for protective action against induced experimental liver tumors in male Wistar rats. The levels of mitochondrial lipid peroxidation (LPO) and liver weight were found to be decreased by the administration of extract (200 and 400 mg/kg) in dose dependent manner. The extract also increased the levels of mitochondrial enzymatic antioxidants²³.

Respiratory and gastrointestinal activity

Methanolic extract from the stem bark was tested for spasmolytic, bronchodilator, and vasodilator activities. The observations confirmed the bronchodilator and vasodilator activities which were possibly mediated through blockade of Ca²⁺ channels²³.

Anticonvulsant activity

Methanolic extract of stem barks was studied for anticonvulsant activity against maximal electro shock (MES) and Pentylenetetrazole (PTZ) induced convulsions in mice. Methanolic extract of stem

barks showed significant anticonvulsant effect in both models²³.

References

1. Goyal M, Sharma SK. Traditional wisdom and value addition prospects of arid foods of desert region of North West India. *Indian Journal of Traditional Knowledge*. 8, 2009, 581-585.
2. Sharma Alok, Shanker C., Tyagi Lalit Kumar, Singh Mahendra and Rao Ch.V., Herbal Medicine for Market Potential in India: An Overview. *Acad J Plant Sci*. 2008;1 (2): 26.
3. Duraipandiyar V, Ayyanar M, Ignacimuthu S, Antimicrobial Activity of Some Ethnomedical Plants Used by Paliyar Tribe from Tamil Nadu, India. *BMC complementary and alternative medicine*. 2006; 635.
4. Puri S. and Kumar A., Establishment of *Prosopis cineraria* (L.) Druce in the hot deserts of India. *New forests*. 1995; 9: 21-33.
5. Sass, J.E. *Elements of Botanical Micro Technique*, McGraw Hill Book Co; New York; 1940.p.222
6. Mahoney D., *Trees of Somalia - A field guide for development workers*, Oxfam/HDRA, Oxford. (1990):133-136.
7. Kirtikar K.R. and Basu B.D. *Indian medicinal plants*. Vol. II. International Book Distributors, Dehradun, India. (1984): 910.
8. Orwa C, Mutua A, Kindt R, Jamnadass R, Simons A. 2009. *Agroforestry Database: a tree reference and selection guide version 4.0* (<http://www.worldagroforestry.org/af/treedb/>).
9. Rastogi RP and BN Mehrotra. *Compendium of Indian Medicinal Plants*. A CDRI Series (Vol IV) Lukhnow: Publication & Information Directorate, New Delhi, 1995.
10. Gangal S., Sharma S and Rauf A. Fatty Acid Composition of *P. cineraria* seeds, *Chem. Nat Compounds*. 2009; 45(5); 705-707.
11. Khan ST, Riaz N, Afza N, et al. Studies on the chemical constituents of *Prosopis cineraria*. *Journal of the Chemical Society of Pakistan*. 2006;28(6):619-622.
12. Malik A, Kalidhar S. Phytochemical examination of *Prosopis cineraria* L.(druce) leaves. *Indian Journal of Pharmaceutical Sciences*. 2007; 69 (4):576-578.
13. Malik A, Khalidar SB. Phytochemical examination of *Prosopis cineraria* leaves. *Indian Journal of Pharmaceutical Sciences*. 2007; 69: 576-578.
14. Ruchika Sharma*, Nandini Jodhawat, Sanju Purohit and Swarnjeet Kaur. Antibacterial activity and Phytochemical screening of *Prosopis Cineraria*. *Int. J. Pharm. Sci. Rev. Res.*, 14(1), 2012; 04, 15-17.
15. Purohit SD, Ramawat KG, Arya HC. Phenolics, peroxidase and phenolase as related to gall formation in some arid zone plants. *Curr Sci* 1979; 48: 714-16.
16. Rhoades DF. *Herbivores, their interaction with secondary plant metabolites*. Acad Press Inc London 1979: pp 3-54.
17. N.A.S., *Firewood crops*. Shrub and tree species for energy production. National Academy of Sciences, Washington D C, 1980a.
18. ICFRE, (Indian Council of Forestry Research and Education), Khejri (*Prosopis cineraria*) ICFRE, Dehradun, India, 1993.
19. Shalini. *Vedic Leguminous Plants*, (Shalini ed) 1997: pp57-8.
20. Toky OP. *Medicinal values of Prosopis cineraria in arid and semiarid India*. Society of chemical industry, I, 1999.
21. Joseph A, George H, Sharma A. & Gopal N. Antipyretic and Analgesic Effects of the Aqueous extract of *P.cineraria*. *Global Journal Of Pharmacology*, 2011,5(2): 73-77.
22. Sharma N, Garg V and Paul Arpita. Antihyperglycemic, Antihyperlipidemic And Antioxidative Potential Of *Prosopis Cineraria* Bark, *Indian Journal of Clinical Biochemistry*, 2010 / 25 (2) 193-200.
23. Vijay N, Sumitra S, Surendra S. Antibacterial Activity of Stem Bark of *Salvadora Oleoides* Decne. *International Journal of Pharmacognosy and Phytochemical Research*. 2013; 5:1: 76-78.